

SYNERGISTIC FUNGICIDAL COMPOSITION COMPRISING A
COMPOUND ANALOGOUS TO STROBILURIN

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation of Application No. 09/505,874, filed February 17, 2000, incorporated by reference herein in its entirety and relied upon, which is a divisional of Application No. 09/051,199, filed July 2, 1998, now U.S. Patent No. 6,057,331, which is the U.S. national phase of PCT/FR96/01532, filed October 2, 1996, and designating the United States, and published by the International Bureau in French, not English, on April 10, 1997 as WO97/12520.

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

[0002] The subject of the present invention is a synergistic fungicidal composition comprising a compound analogous to strobilurin and a process which makes use of the said composition and which is intended for the curative or preventive protection of crops against fungal attacks.

BACKGROUND OF THE PRIOR ART

[0003] Compounds analogous to strobilurin with a fungicidal effect are known, in particular from European

Patent Publication EP 253213 or 398692 or from International Publication WO 9208703, which make it possible to prevent the growth and the development of phytopathogenic fungi capable of attacking crops.

[0004] However, it is always desirable to improve the spectrum of activity and the effectiveness of such compounds with a fungicidal effect or to reinforce them by combining them with other molecules in order to obtain a more effective product or alternatively to prevent the appearance of fungal strains which are resistant to these new fungicides.


[0005] It is also highly desirable to have available fungicidal products which enjoy an improved persistence of effect, likely to space out in time the number of plant-protection treatments necessary for satisfactory control of the parasites.

[0006] It is, in any event, particularly advantageous to be able to decrease the amount of chemicals distributed in the environment, while ensuring high-performance protection of crops against fungal attacks.

[0007] It has now been found that one (or a number) of the above objectives could be achieved by virtue of the fungicidal composition according to the present invention.

SUMMARY OF THE INVENTION


[0008] The subject of the present invention is therefore a synergistic fungicidal composition comprising at least one compound A of formula (I):



in which:

A is the nitrogen atom or the -CH group,

B is the -OCH₂-, -CH₂O-, -CH(CH₃)-O-N=CH- or -CH=N-O-CH(CH₃)- group or the group of formula (II) :



R₁ is an alkyl group containing 1 to 4 carbon atoms, preferably the methyl group,

R₂ is -OCH₃ or -NHCH₃,

X is a halogen atom, the cyano group or an alkyl or haloalkyl group containing 1 to 4 carbon atoms, preferably the methyl or trifluoromethyl group,

n is equal to 1 or 2, with, when n is equal to 2, the possibility of having different X groups,

and at least one fungicidal compound B chosen from the group comprising dicarboximide derivatives, such as iprodione, procymidone and vinchlozolin,

the said composition comprising the components A and B in an A/B ratio by weight of between 0.02 and 5, preferably between 0.0625 and 1.33,

with the proviso that when compound B is procymidone, then B in formula I is the group of formula (II).

BRIEF DESCRIPTION OF THE DRAWING

[0009] The sole Figure of Drawing, FIGURE 1, is an isobole plot for ICIA 5504, i.e. methyl (E)-2-{2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate, and iprodione in radish seedlings against *Alternaria brassicae* (alternaria disease).

DETAILED DESCRIPTION OF THE INVENTION, PREFERRED EMBODIMENTS AND BEST MODE

[0010] It is clearly understood that the said fungicidal composition can contain a single compound B or more than one such compound, for example 1, 2 or 3 compounds B, according to the use for which the

combination is intended. Likewise, the composition may comprise more than one compound A.

[0011] Preference is given to the fungicidal composition according to the invention in which the compound A is methyl (E)-2-{2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate, methyl (E)-methoxyimino[α -(o-tolyloxy)-o-tolyl]acetate or N-methyl-(E)-methoxyimino[2-(2,5-dimethylphenoxyethyl)phenyl]acetamide.

[0012] Preference is further given, among the more especially preferred meanings of the compound B defined above, to iprodione. In an entirely unexpected way, the composition according to the invention significantly improves the effect of the active materials taken separately with respect to a certain number of fungi which are particularly harmful to crops, for example in particular grapes or the Solanaceae. This improvement is reflected in particular by a decrease in the doses of each of the constituents, which is particularly advantageous for the user and the environment. The fungicidal product thus exhibits synergistic properties attested by application of the Tamme's method, "Isoboles, a graphic representation of synergism in pesticides", Netherlands Journal of Plant Pathology, 70(1964), p. 73-80 or as defined by Limpel, L.E., P.H. Schuldt and D. Lammont, 1962, Proc. NEWCC 16:48-53, using the following formula, also called Colby's formula:

$$E = X + Y - X.Y/100$$

in which:

- E is the expected percentage inhibition of the growth of the fungus by a mixture of the two fungicides A and B at defined doses, equal to a and b respectively;

- X is the observed percentage inhibition by the fungicide A at dose a,

- Y is the observed percentage inhibition by the fungicide B at dose b.

[0013] When the percentage inhibition observed for the mixture is greater than E, there is synergism.

[0014] Preferably, when the component B is iprodione, the A/B ratio is between 0.05 and 5, preferably between 0.17 and 1.33, for all the crops envisaged.

[0015] In the particular case of lawns, the A/B ratio will generally be between 0.02 and 2, preferably between 0.0625 and 0.25.

[0016] The structures corresponding to the common names of the fungicidal active materials which appear in the definition of B are shown in at least one of the following 2 works:

- "The Pesticide Manual", edited by Clive Tomlin and published by The British Crop Protection Council, 10th edition;

- l'Index phytosanitaire [Plant-protection index] 1994, published by l'Association de Coordination Technique Agricole [Agricultural Technical Coordination Association], 30th edition.

200207484001

[0017] As regards the derivatives analogous to strobilurin, methyl (E)-2-{2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate (or ICIA5504) is described in International Publication WO 9208703; methyl (E)-methoxyimino[α -(o-tolyloxy)-o-tolyl]acetate (or BAS490F) is described in European Patent Publication EP 253213; and N-methyl-(E)-methoxyimino[2-(2,5-dimethylphenoxy)methyl]phenyl]acetamide (or SSF-129) is described in European Patent Publication EP 398692. Moreover, ICIA5504 and BAS490F are listed in the abovementioned work "The Pesticide Manual".

[0018] The fungicidal composition according to the invention comprises, as active material, at least one compound A and at least one compound B as a mixture with solid or liquid vehicles which are acceptable in agriculture and/or surface-active agents which are also acceptable in agriculture. In particular, the inert and conventional vehicles and the conventional surface-active agents can be used. These compositions cover not only compositions which are ready to be applied to the crop to be treated by means of a suitable device, such as a spray device, but also commercial concentrated compositions which have to be diluted before application to the crop. The combination of at least one compound A with at least one compound B is denoted as active material.

[0022] The surface-active agent can be an emulsifying, dispersing or wetting agent of ionic or nonionic type or a mixture of such surface-active agents. There may be mentioned, for example, salts of polyacrylic acids, salts of lignosulphonic acids, salts of phenolsulphonic or naphthalenesulphonic acids, polycondensates of

ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols, (especially alkylphenols or arylphenols), salts of esters of sulphosuccinic acids, taurine derivatives (especially alkyltaurates), phosphoric esters of polyoxyethylenated phenols or alcohols, esters of fatty acids and of polyols, and the derivatives of the above compounds containing sulphate, sulphonate and phosphate functional groups. The presence of at least one surface-active agent is generally indispensable when the active material and/or the inert vehicle is/are not soluble in water and the carrier agent for application is water.

[0023] The compositions for agricultural use according to the invention can thus contain the active material within very wide limits, ranging from 0.05 % to 95 % (by weight). Their surface-active agent content is advantageously of between 5 % and 40 % by weight. Unless otherwise stated, the percentages given in this description, including the claims, are by weight.

[0024] These compositions according to the invention are themselves in fairly diverse, solid or liquid forms.

[0025] There may be mentioned, as forms of solid compositions, the powders for dusting (with an active material content which can range up to 100 %) and the granules, especially those obtained by extrusion, by compacting, by impregnation of a granulated support, or by granulation from a powder (the active material

content in these granules being between 0.5 and 80 % for the latter cases), the tablets or effervescent tablets.

[0026] The fungicidal composition according to the invention can also be used in the form of powders for dusting; it is also possible to use a composition comprising 50 g of active material and 950 g of talc; it is also possible to use a composition comprising 20 g of active material, 10 g of finely divided silica and 970 g of talc; these constituents are mixed and milled and the mixture is applied by dusting.

[0027] There may be mentioned, as forms of liquid compositions or those intended to constitute liquid compositions at the time of application, solutions, in particular water-soluble concentrates, emulsifiable concentrates, emulsions, suspension concentrates, aerosols, wettable powders (or powder to be sprayed), pastes or gels.

[0028] The emulsifiable or soluble concentrates most often comprise 10 to 80 % of active material, the emulsions or solutions ready for application containing, for their part, 0.001 to 20 % of active material.

[0029] In addition to the solvent, the emulsifiable concentrate may contain, when necessary, 2 to 20 % of appropriate additives such as stabilizing agents, surface-active agents, penetrating agents, corrosion inhibitors, colorants or adhesives mentioned above.

[0030] Starting from these concentrates, it is possible to obtain, by diluting with water, emulsions of any

desired concentration which are particularly suitable for applying to crops.

[0031] By way of example, here is the composition of a few emulsifiable concentrates:

EC EXAMPLE 1:

- active material	400 g/l
- alkali metal dodecylbenzene sulphonate	24 g/l
- oxyethylated nonylphenol with 10 molecules of ethylene oxide	16 g/l
- cyclohexanone	200 g/l
- aromatic solvent	qs 1 litre

[0032] According to another emulsifiable concentrate formula, the following are used:

EC EXAMPLE 2:

- active material	250 g
- expoxidized vegetable oil	25 g
- mixture of alkylaryl sulphonate and of ether of polyglycol and fatty alcohols	100 g
- dimethylformamide	50 g
- xylene	575 g

[0033] The suspension concentrates, which can also be applied by spraying, are prepared so as to obtain a stable fluid product which does not settle out and they

generally contain from 10 to 75 % of active material, from 0.5 to 15 % of surface-active agents, from 0.1 to 10 % of thixotropic agents and from 0 to 10 % of suitable additives, such as antifoaming agents, corrosion inhibitors, stabilizing agents, penetrating agents and adhesives and, as vehicle, water or an organic liquid in which the active material is insoluble or nearly insoluble: certain organic solid materials or inorganic salts can be dissolved in the vehicle to aid in preventing sedimentation or as antigels for water.

[0034] A suspension concentrate composition is given here as an example:

SC EXAMPLE 1:

- active material	500 g
- polyethoxylated tristyrylphenyl phosphate	50 g
- polyethoxylated alkylphenol	50 g
- sodium polycarboxylate	20 g
- ethylene glycol	50 g
- organopolysiloxane oil (antifoaming agent)	1 g
- polysaccharide	1.5 g
- water	316.5 g

[0035] Wettable powders (or powder to be sprayed) are generally prepared so that they contain 20 to 95 % of active material, and they generally contain, in addition to the solid vehicle, from 0 to 30 % of a wetting agent, from 3 to 20 % of a dispersing agent and, when this is

necessary, from 0.1 to 10 % of one or more stabilizing agents and/or other additives, such as penetrating agents, adhesives, or anticlumping agents, dyes, and the like.

[0036] To obtain powders to be sprayed or wettable powders, the active materials are intimately mixed, in suitable mixers, with the additional substances and the mixture is milled with mills or other suitable grinders. Powders to be sprayed are thereby obtained with advantageous wettability and suspensibility; they can be suspended in water at any desired concentration and these suspensions can be used very advantageously in particular for application to plant leaves.

[0037] Pastes can be produced in place of wettable powders. The conditions and modes of production and use of these pastes are similar to those of wettable powders or powders to be sprayed.

[0038] Various wettable powder (or powder to be sprayed) compositions are given here as examples:

WP EXAMPLE 1

- active material	50 %
- ethoxylated fatty alcohol (wetting agent)	2.5 %
- ethoxylated phenylethylphenol (dispersing agent)	5 %
- chalk (inert vehicle)	42.5 %

WP EXAMPLE 2:

- active material 10 %
- C13 branched-type synthetic oxo alcohol,
ethoxylated with 8 to 10 molecules of
ethylene oxide (wetting agent) 0.75 %
- neutral calcium lignosulphonate
dispersing agent) 12 %
- calcium carbonate (inert filler) qs for 100 %

[0039] This wettable powder contains the same ingredients as in the above example, in the proportions below:

WP EXAMPLE 3:

- active material 75 %
- wetting agent 1.50 %
- dispersing agent 8 %
- calcium carbonate (inert filler) qs for 100 %

WP EXAMPLE 4:

- active material 90 %
- ethoxylated fatty alcohol (wetting agent) 4 %
- ethoxylated phenylethylphenol
(dispersing agent) 6 %

WP EXAMPLE 5:

- active material 50 %
- mixture of anionic and nonionic surface-active agents (wetting agent) 2.5 %
- sodium lignosulphonate (dispersing agent) 5 %
- kaolin clay (inert vehicle) 42.5 %

[0040] Aqueous dispersions and emulsions, for example the compositions obtained by diluting a wettable powder or an emulsifiable concentrate according to the invention with water, are contained within the general scope of the present invention. Emulsions can be of the water-in-oil or oil-in-water type and they can have a thick consistency, like that of a "mayonnaise".

[0041] The fungicidal compositions according to the invention can be formulated in the form of water-dispersible granules, which also come within the scope of the invention.

[0042] These dispersible granules, with a bulk density generally of between approximately 0.3 and 0.6, have a particle size generally of between approximately 150 and 2000 and preferably between 300 and 1500 microns.

[0043] The active material content of these granules is generally between approximately 1 % and 90 %, and preferably between 25 % and 90 %.

[0044] The rest of the granule comprises a solid filler and, optionally, surface-active adjuvants which confer water-dispersibility properties on the granule. These granules can be essentially of two distinct types

according to whether the filler chosen is soluble or insoluble in water. When the filler is water-soluble, it can be inorganic or, preferably, organic. Excellent results were obtained with urea. In the case of an insoluble filler, the latter is preferably inorganic, such as, for example, kaolin or bentonite. It is then advantageously accompanied by surface-active agents (in a proportion of 2 to 20 % by weight of the granule) of which more than half comprises, for example, at least one dispersing agent, essentially anionic, such as an alkali metal or alkaline-earth metal polynaphthalenesulphonate or an alkali metal or alkaline-earth metal lignosulphonate, the remainder consisting of nonionic or anionic wetting agents such as an alkali metal or alkaline-earth metal alkylnaphthalenesulphonate.

[0045] Moreover, although this is not indispensable, it is possible to add other adjuvants such as antifoaming agents.

[0046] The granule according to the invention can be prepared by mixing the necessary ingredients and then granulating according to several techniques known per se (granulator, fluid bed, sprayer, extrusion, and the like). The preparation generally finishes with a crushing followed by a sieving to the particle size chosen within the limits mentioned above. Granules obtained as above and then impregnated with a composition containing the active material can alternatively be used.

[0047] It is preferably obtained by extrusion, by carrying out the preparation as indicated in the examples below.

DG EXAMPLE 1: DISPERSIBLE GRANULES

[0048] 90 % by weight of active material and 10 % of urea in the form of pearls are mixed in a mixer. The mixture is then milled in a pin mill. A powder is obtained which is moistened with approximately 8 % by weight of water. The moist powder is extruded in a perforated-roller extruder. A granular material is obtained which is dried, and then crushed and sieved, so as to respectively keep only the granules with a size of between 150 and 2000 microns.

DG EXAMPLE 2: DISPERSIBLE GRANULES

[0049] The following constituents are mixed in a mixer:

- | | |
|---|------|
| - active material | 75 % |
| - wetting agent (sodium alkylnaphthalene-sulphonate | 2 % |
| - dispersing agent (sodium polynaphthalenesulphonate) | 8 % |
| - water-insoluble inert filler (kaolin) | 15 % |

[0050] This mixture is granulated on a fluid bed in the presence of water and then dried, crushed and sieved so as to obtain granules with a size of between 0.15 and 0.80 mm.

[0055] More precisely, the subject of the invention is a process for the curative or preventive control of fungi which are phytopathogenic, for example, towards crops or lawns, characterized in that an effective and non-phytotoxic amount of a combination of at least one compound A and at least one compound B, for example in a fungicidal composition according to the invention, is generally applied on the aerial parts of plants.

- from the Oomycetes group:
 - from the *Phytophthora* genus, such as *Phytophthora infestans* (downy mildew of the Solanaceae, in particular of potatoes or tomatoes), *Phytophthora citrophthora*, *Phytophthora capsici*, *Phytophthora cactorum*, *Phytophthora palmivora*, *Phytophthora cinnamoni*, *Phytophthora megasperma* or *Phytophthora parasitica*,
 - from the Peronosporaceae family, in particular *Plasmopara viticola* (grape downy mildew), *Plasmopara halstedei* (sunflower downy mildew), *Pseudoperonospora* sp (in particular downy mildew of cucurbits and of hops), *Bremia lactucae* (lettuce downy mildew) or *Peronospora tabacinae* (tobacco downy mildew),
 - from the Adelomycetes group:
 - from the *Alternaria* genus, for example *Alternaria solani* (alternaria disease of the Solanaceae and in particular of tomatoes and potatoes),
 - from the *Guignardia* genus, in particular *Guignardia bidwellii* (black rot of grapes),
 - from the *Oidium*s group, for example grape powdery mildew (*Uncinula necator*); powdery mildew

of legumes, for example *Erysiphe polygoni* (powdery mildew of crucifers); *Leveillula taurica*, *Erysiphe cichoracearum*, *Sphaerotheca fuligena*; (powdery mildew of cucurbits, of composites or of tomatoes); *Erysiphe communis* (beet and cabbage powdery mildew); *Erysiphe pisi* (pea or lucerne powdery mildew); *Erysiphe polyphaga* (bean and cucumber powdery mildew); *Erysiphe umbelliferarum* (powdery mildew of the Umbelliferae, in particular of carrots); *Sphaerotheca humuli* (hop powdery mildew); or *Erysiphe graminis* (powdery mildew of cereals),

- from the *Septoria* genus, for example *Septoria nodorum* or *Septoria tritici* (septoria disease of cereals);

- from the *Sclerotinia* genus, for example *Sclerotinia sclerotinium* or *Sclerotinia homeocarpa* (lawn sclerotinia disease),

- from the Basidiomycetes group:

- from the *Puccinia* genus, for example *Puccinia recondita* or *striiformis* (wheat rusts),

- of the *Rhizoctonia* spp. family.

[0057] A classification made no longer by targeted fungi but by target crops can be illustrated as below:

- barley-helminthosporium
(*Helminthosporium*),

2007/04/14 14:00:00

- grapes: powdery mildew (*Uncinula necator*), downy mildew (*Plasmopara viticola*), rot (*Botrytis cinerea*), excoriosis (*Phomopsis viticola*) and black rot (*Guignardia bidwellii*),

- legumes: downy mildews (*Peronospora* spp., *Bremia lactucae*, *Pseudoperonospora* spp.), or alternaria disease (*Alternaria* spp.), sclerotinia disease (*Sclerotinia* spp.), rot (*Botrytis cinerea*) or powdery mildew (*Erysiphe* spp. or *Sphaerotheca fuliginea*), foot or root rot (*Rhizoctonia* spp.),

- rice: foot rot or root rot (*Rhizoctonia* spp.),

- arboriculture: scab (*Venturia*
inaequalis), powdery mildew (*Podosphaera leucotricha*),
alternaria disease (*Alternaria* spp.), rot (*Botrytis*
cinerea) and monilia disease (*Monilia fructigena*),

- citrus fruits: scab (*Elsinoe fawcetti*),
melanose (*Phomopsis citri*) and diseases due to
Phytophthora spp.,

bananas: cercospora disease
(*Mycosphaerella figiensis*),

- lawns: rust, powdery mildew,
helminthosporium disease or soil diseases (*Microdochium*
nivale, *Pythium* spp., *Rhizoctonia solani*, *Sclerotinia*
homeocarpa and the like).

[0058] The fungicidal composition which is the subject
of the invention is applied by means of various
treatment processes, such as:

- spraying a liquid comprising the said
composition on the aerial parts of the crops to be
treated,

- dusting, the incorporation of granules or of
powders into the soil, sprinkling, injecting into trees
or painting.

[0059] The spraying of a liquid on the aerial parts of
the crops to be treated is the preferred treatment
process.

[0060] "Effective and non-phytotoxic amount" is
understood to mean an amount of composition according to
the invention which is sufficient to make possible
control or destruction of the fungi present or capable
of appearing on the crops and which does not result in
any notable phytotoxicity symptoms with respect to the

[0061] The use doses during the implementation of the process according to the invention will then generally be:

100 to 1000 g of compound B, e.g. iprodione
+ 50 to 500 g/ha of compound A and more precisely 300 to
600 g + 100 to 400 g/ha, i.e. a total dose of compounds
A and B of between 150 and 1500 g/ha, preferably between
400 and 1000 g/ha,

1000 to 5000 g of compound B, e.g. iprodione
+ 100 to 2000 g/ha of compound A and more precisely 2000
to 4000 g + 250 to 1000 g/ha, i.e. a total dose of
compounds A and B of between 1100 and 7000 g/ha,
preferably between 2250 and 5000 g/ha.

[0063] In the figure appended to the present text, the dose of each active material, taken in isolation,

required for control of the phytopathogenic fungus at the level indicated is compared with that of the 2 active materials taken as a mixture. The effective dose of each active material taken in isolation is shown on the axis of the abscissae and of the ordinates and a straight line is traced cutting these 2 axes and connecting these 2 doses. When an active material taken in isolation is not effective, the straight line is parallel to the axis of the coordinates which shows the doses of this active material. As regards the 2 active materials taken as a mixture, the dose of the mixture at a given ratio is shown by a point.

EXAMPLE 1: In vivo test of the combination of ICIA5504 with iprodione with respect to alternaria disease of radish (*Alternaria brassicae*) by preventive treatment 24 hours before infection.

[0064] A suspension comprising the compounds A and B in a liquid mixture composed of a surface-active agent (oleate of polyoxyethylenated derivative of sorbitan) and water is prepared.

[0065] The component B is iprodione; the A/B ratio is 0.25, 0.5, 1, 2 or 4.

[0066] Radish seedlings (variety Pernot), are grown in small pots. When these seedlings are 10-12 days old (cotyledon, 1st leaf stage), they are treated by spraying with the above suspension.

[0067] Seedlings, used as controls, are treated with a similar suspension but which does not contain active material ("formulation blank").

[0068] After 24 hours, each seedling is infected by spraying with an aqueous suspension of spores of *Alternaria brassica*, which suspension is obtained from an in vitro culture of the fungus. These spores are suspended at a concentration of 40,000 units per cm³ of inoculum. Infection is carried out by spraying the inoculum at the top surface of the leaves.

[0069] The infected seedlings are then incubated for six days at 18-20°C at 90-100 % relative humidity under natural light.

[0070] Reading is carried out 6 days after infection, in comparison with the control seedlings.

[0071] The results obtained are reported in the form of points, corresponding to 80% destruction of the parasite, and are plotted on a Tamme isobole diagram which contains, on the abscissa, the doses of A expressed in ppm (mg/l) and, on the ordinate, the doses of B, also in ppm (mg/l).

[0072] The diagram of Figure 1 is obtained. It appears that the addition of iprodione makes it possible, in an entirely unexpected way, to lower the dose of A necessary for the destruction of 80 % of the parasite below 115 ppm (mg/l), which corresponds to the dose of A alone which it is necessary to apply to obtain the same percentage of destruction.

[0073] The arrangement of the points obtained therefore shows a bilateral effect described, according to the Tammes method mentioned above, as a "two-sided effect". This arrangement corresponds to a type-III isobole according to the said method (page 74 of the corresponding bibliographical reference already cited) and is characteristic of a synergy.

EXAMPLE 2: In vivo test of the combination of ICIA5504 with iprodione with respect to helminthosporium disease of barley by preventive treatment 7 days before infection.

[0074] The fungicides are used at the following doses:

iprodione (Rovral SC 500 g/l (B): 12.5-25-50-100-200-400-800 g/ha,

ICI A5504 (SC 250 g/l) (A): 1.5-3.1-6.2-12.5-25-50-100-200 g/ha,

iprodione (B)/ICIA5504 (A) = 4: 6.2+1.5 to 800+200 g/ha,

iprodione/ICIA5504 = 2: 3.1+1.5 to 400+200 g/ha

iprodione/ICIA5504 = 1: 1.5+1.5 to 200+200 g/ha

iprodione/ICIA5504 = 0.5: 0.75+1.5 to 100+200 g/ha

[0075] The young barley plants (var. Express) which are 12 days old (first leaf developed stage) are treated with the experimental fungicides at the doses indicated above in a slurry volume of 250 l/ha (3 repeats/dose). Seven days after the treatment, the young barley plants

are infected with a suspension of *Pyrenophora teres* spores (10,000 spores/ml). The young plants are incubated for 48 hours at 20°C, 100 % relative humidity (RH). Next, the plants are transferred to a greenhouse at 20°C, 70-80 % RH under natural light. The grading of the disease is carried out ten days after infection by estimating the leaf surface area contaminated on the first leaf (10 young plants/pot).

[0076] By comparing with the control (non-treated-contaminated) a percentage efficacy is calculated. The synergy is then quantified by using the Colby model. The following results are thereby obtained:

Observed efficacy:

B/A	0 g/ha	1.5	3.1	6.2	12.5	25	50	100	200
0 g/ha	0	0	17.54	34.21	59.21	76.75	83.86	81.67	92.72
12.5	15.79		78.6	88.68	83.95	94.04			
25	7.02			88.77	94.47	85.70	93.68		
50	37.72				91.49	96.23	92.54	90.88	
100	35.09					94.65	95.53	93.33	96.23
200	47.37						95.53	97.37	95
400	50.88							96.14	97.72
800	59.65								95.95

Theoretical efficacy according to Colby:

B/A	0	1.5	3.1	6.2	12.5	25	50	100	200
g/ha	g/ha								
0 g/ha	0	0	17.54	34.21	59.21	76.75	83.86	81.67	92.72
12.5	15.79		30.56	44.60	65.65	80.42			
25	7.02			38.83	62.07	78.38	84.99		
50	37.72				74.60	85.52	89.95	88.58	
100	35.09					84.91	89.52	88.10	95.27
200	47.37						91.51	90.35	96.17
400	50.88							91.00	96.42
800	59.65								97.06

